

Unclassified

OTIC FILE COPY

(4)

SECURITY CLASSIFICATION OF THIS PAGE

PORT DOCUMENTATION PAGE

1a. AD-A200 926			1b. RESTRICTIVE MARKINGS	
2a. IC			3. DISTRIBUTION/AVAILABILITY OF REPORT Distribution List Enclosed	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NOV 09 1988			5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6a. NAME OF PERFORMING ORGANIZATION Purdue University		6b. OFFICE SYMBOL (If applicable)	7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Arlington, Virginia 22217	
6c. ADDRESS (City, State, and ZIP Code) West Lafayette, Indiana 47907			9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-86-K-0547	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)	10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217-5000			PROGRAM ELEMENT NO.	PROJECT NO.
			TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Physical and Chemical Characteristics of Organoboranes				
12. PERSONAL AUTHOR(S) Herbert C. Brown				
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM 8/6/86 TO 9/30/88		14. DATE OF REPORT (Year, Month, Day) 1988, October 19
15. PAGE COUNT 5				
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	Ipc ₂ BX, meso-epoxides, enantioselective ring cleavage, 1,2-halohydrins, R ₂ BCl, R ₂ BOTf, enol borinates, hydroboration	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) see attached report				
<div style="border: 1px solid black; padding: 5px; text-align: center;"> DISTRIBUTION STATEMENT A Approved for public release Distribution Unlimited </div>				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> OTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Professor Herbert C. Brown			22b. TELEPHONE (Include Area Code) (317) 494-5316	22c. OFFICE SYMBOL

OFFICE OF NAVAL RESEARCH

FINAL REPORT

for

Contract N00014-86-K-0547

Physical and Chemical Characteristics of Organoboranes



Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By <i>per ltr</i>	
Distribution	
Availability Codes	
Dist	For and/or Special
A-1	

Herbert C. Brown
Department of Chemistry
Purdue University
West Lafayette, IN 47907

October 19, 1988

88 1025 017

Personnel

R. K. Dhar
S. Swaminathan
P. Pandiarajan
C. Zhao
M. Srebnik
G. Rajendran

Period on the Budget

10/1/86 - 9/30/88
8/5/86 - 6/30/88
12/1/86 - 5/31/87
10/1/87 - 10/18/88
1/1/88 - 9/30/88
2/1/88 - 7/31/88

Technical Reports and Journal Articles**Technical Reports**

1. Enantioselective Ring-Cleavage of *meso*-Epoxides with B-Halodiisopinocampheylboranes
N. N. Joshi, M. Srebnik and H. C. Brown
2. A Comparative Study of Dialkylboron Chlorides and Triflates for the Enolization of Ketones.
The Controlled Stereospecific Synthesis of Either [*E*]- or [*Z*]-Enol Borinates
H. C. Brown, B. Singaram, R. K. Dhar, P. K. Pandiarajan and R. K. Dhar
3. Enantioselective Ring-Cleavage of *meso*-Epoxides with B-Halodiisopinocampheylboranes
N. N. Joshi, M. Srebnik and H. C. Brown
Oral presentation at the 146th Meeting of the American Chemical Society in Los Angeles,
CA, September 25-30, 1988

Journal Articles


1. Enantioselective Ring-Cleavage of *meso*-Epoxides with B-Halodiisopinocampheylboranes
N. N. Joshi, M. Srebnik and H. C. Brown
J. Am. Chem. Soc. **1988**, *110*, 6246

Summary of Accomplishments**1. Enantioselective Ring Cleavage**

This subject has been detailed in Technical Reports (1), April 26, 1988, and (3), May 18, 1988. We have succeeded in obtaining optically active 1,2-halohydrins from *meso*-epoxides and chiral organoboranes. This is the first time such an asymmetric transformation has been accomplished.


epoxides. (1988) ←

①



2. Enol Borinates

This topic was the subject of Technical Report (2), May 13, 1988. In essence we have found conditions for obtaining [*E*]-enol borinates from ketones and readily available dialkylchloroboranes.



Reaction of the [*E*]-enol borinate with an aldehyde then gives the *anti*-aldol product. This methodology therefore complements the known process of obtaining [*Z*]-enol borinates from ketones and dialkyl boron triflates, which in turn provide *syn*-aldol products.





3. Hydroboration of Alkenes

This project was only briefly covered in the End-of-the-Year Reports (July 21, 1987 and July 22, 1988) and therefore will now be presented in greater detail.

Monohydroboration of symmetrically substituted internal alkynes, such as 3-hexyne and diphenylacetylene, with an equivalent of 9-BBN can be carried out at 0°C and 25°C respectively, producing the B-vinyl derivatives in 95-98% yields. The dihydroboration of 3-hexyne (0.25 *M*) with 2 equiv of 9-BBN (0.5 *M*) can only produce 56% of *gem*-dibora derivatives with 44% of B-vinyl derivatives at 25°C. However, the formation of *gem*-dibora derivatives can be improved by increasing the temperature (65°C) and concentration (0.75 *M*). The reaction of diphenylacetylene with 2 equiv of 9-BBN can only produce B-vinyl derivatives at 25°C, with no dihydroborated product. The dihydroboration of terminal alkynes, such as 1-hexyne and phenylacetylene, with 2 equiv of 9-BBN can produce *gem*-dibora derivatives in 81-90% yields. But the monohydroboration of 1-hexyne and phenylacetylene with an equivalent of 9-BBN can only

provide a mixture of B-vinyl and *gem*-dibora derivatives in 64:36 and 76:24 respectively. Fortunately, the monohydroboration of 1-hexyne with 9-BBN can be improved by use of 100% of excess of terminal alkyne, making possible the preparation of the corresponding B-[E]-1-hexenyl-9-BBN in 85% yield with no dihydroborated product. But in the case of phenylacetylene, the monohydroboration can only be improved to 83:17 from 76:24.

No general synthesis of B-vinyl and dibora derivatives has been studied in detail using various hydroborating agents and alkynes. Consequently, a systematic study was undertaken to prepare various B-alkenyl and dibora derivatives from representative alkynes using 9-BBN (eqs 1-4).



All yields were $\geq 80\%$. The products were confirmed by conversion to the corresponding aldehydes.